

## Original article

## Asian Pacific Journal of Tropical Medicine

journal homepage: [www.apjtm.org](http://www.apjtm.org)

doi: 10.4103/1995-7645.254941

Impact factor: 1.634

## Occurrence of *Escherichia coli* virulence genes in feces of wild birds from Central Italy

Fabrizio Bertelloni<sup>1</sup>, Errica Lunardo<sup>1</sup>, Guido Rocchigiani<sup>1</sup>, Renato Ceccherelli<sup>2</sup>, Valentina Virginia Ebani<sup>1</sup>✉

<sup>1</sup>Department of Veterinary Science, University of Pisa, viale delle Piagge 2, 56124 Pisa, Italy

<sup>2</sup>CRUMA-LIPU, via delle Sorgenti 430, 57121 Livorno, Italy

## ARTICLE INFO

## Article history:

Received 5 November 2018

Received in revised form 26 January 2019

Accepted 15 February 2019

Available online 27 March 2019

## Keywords:

Wild birds

Enteropathogenic *Escherichia coli*Enterohemorrhagic *Escherichia coli*Shiga-toxins producing *Escherichia coli*

## ABSTRACT

**Objective:** To investigate the potential role of wild birds as fecal spreaders of enteropathogenic, enterohemorrhagic and Shiga-toxins producing *Escherichia coli* (*E. coli*), enteropathogenic *E. coli*, enterohemorrhagic *E. coli* and Shiga toxin-producing *E. coli* strains.

**Methods:** Fecal samples collected from 121 wild birds of different orders and species were submitted to molecular analyses. In particular, *eaeA* encoding intimin, *hlyA* encoding for hemolysin, *stx1* and *stx2* genes encoding Shiga-toxins 1 and 2, respectively, were investigated.

**Results:** Overall, 21(17.35%) fecal samples resulted positive for at least one of the investigated genes. In detail, 12(9.91%) samples were positive for *eaeA*, 10(8.26%) for *stx1*, 4(3.31%) for *hlyA* and 1(0.83%) for *stx2*. An owl (*Athene noctua*) positive for the four investigated genes suggesting that it harbored a STEC strain. However, virulence genes characterizing EPEC, and EHEC strains were mainly found among seagulls, waterfowl and feral pigeons.

**Conclusions:** Seagulls, waterfowl and feral pigeons, which frequently reach and contaminate rural, urban and peri-urban areas with their droppings, may be important sources of *E. coli* infection for other animals and humans.

### 1. Introduction

*Escherichia coli* (*E. coli*) is an opportunistic Gram negative bacterium presenting in the normal gut microflora of human and animals, even though some strains have been related to a wide range of diseases of veterinary and human concern[1].

Enterohemorrhagic *E. coli* (EHEC) strains are responsible for gastrointestinal disease in humans including diarrhea, hemorrhagic colitis and hemolytic uremic syndrome (HUS), which is the main cause of acute renal failure in children[2]. One of the most important

virulence factors of EHEC is hemolysin, which contributes to the pathogenesis by different mechanisms such as adhesion, induction of pro-inflammatory reactions and epithelial and endothelial cell damage[3].

Some EHEC strains produce Shiga-toxins *Stx1* and *Stx2*, which represent virulence factors causing intestinal cell death and consequent diarrhea[4]. These strains, named Shiga toxin-producing

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: [reprints@medknow.com](mailto:reprints@medknow.com)

©2019 Asian Pacific Journal of Tropical Medicine Produced by Wolters Kluwer- Medknow. All rights reserved.

**How to cite this article:** Bertelloni F, Lunardo E, Rocchigiani G, Ceccherelli R, Ebani VV. Occurrence of *Escherichia coli* virulence genes in feces of wild birds from Central Italy. Asian Pac J Trop Med 2019; 12(3): 142-146.

First author: Fabrizio Bertelloni, Department of Veterinary Science, University of Pisa, viale delle Piagge 2, 56124 Pisa, Italy.

✉Corresponding author: Dr. Valentina Virginia Ebani, DVM, PhD, Department of Veterinary Science, University of Pisa, Viale delle Piagge 2, 56124 Pisa, Italy.

E-mail: [valentina.virginia.ebani@unipi.it](mailto:valentina.virginia.ebani@unipi.it)

*E. coli* (STEC) or verocytotoxic *E. coli* (VTEC), are important zoonotic agents. Among them, O157:H7 is the most common serotype encountered in hemorrhagic colitis and HUS cases, even though other non-O157 *E. coli* serogroups have been involved in human infections[5].

All EHEC express intimin, constitutes an additional virulence factor mediate the intimate attachment of bacteria to epithelial cells and stimulate mucosal immune responses and intestinal crypt hyperplasia, resulting in the characteristic attaching and effacing (A/E) lesions on intestinal epithelial cells[1,2].

*E. coli* strains producing intimin, but not hemolysin, are termed Enteropathogenic *E. coli* (EPEC) and are related to potentially fatal diarrhea, affecting mainly children in developing countries. Bacterial adherence to enterocytes mediated by intimin induce microvilli loss, which could eventually lead to diarrhea[6].

Domestic ruminants, mainly cattle, are considered the main reservoirs of STEC, even though these strains have been found among several domestic and wild mammals' species, too[7].

Following the first report of STEC infection in wild birds in 1997[8], some studies have been carried out to verify the potential role of birds, including poultry, as spreaders of pathogenic *E. coli* worldwide[6,9-17]. Detection of these pathogens in avian population suggests that birds, together with mammals, may be direct or indirect source of infection for humans.

*E. coli* O157:H7 was isolated from layer hens and from pigeons in Italy[18,19]. However, to the best of our knowledge, data about EPEC, EHEC and STEC infections in wild avian population in Italy are not available.

The aim of the present study was to investigate the potential role of wild birds as fecal spreaders of EPEC, EHEC and STEC strains by searching the genes coding for the virulence factors intimin, hemolysin and Shiga-toxins in the feces of free-living birds of several species and from different environments.

## 2. Materials and methods

### 2.1. Animal samples

From January to December 2016, gut samples were collected from 121 free-roaming wild birds. In particular, 56 samples were collected from hunted animals. These birds have been hunted during the hunting season in different wet areas of Central Italy. While the evisceration was performed by the hunters, intestine was collected and sent to the laboratory of the Avian Pathology Section (APS) of the Department of Veterinary Science (University of Pisa), at 4 °C.

Furthermore, 65 samples were collected from birds dead at a bird recovery center located in Central Italy. During the necropsies, a portion of terminal intestine, approximatively from caeca to

cloaca, were collected from each bird and stored at 4 °C until the investigations.

### 2.2. Ethic statement

Regularly hunted and naturally dead birds were used in the study. No birds were sacrificed for the study.

### 2.3. Molecular analyses

DNA was extracted from about 25 mg of each fecal sample with the commercial kit Tissue Genomic DNA Extraction Kit (Fisher Molecular Biology, Trevose, PA, USA), following the manufacturers' guidelines; once obtained, DNA samples were kept at 4 °C until used as template in PCR assays.

Single PCR protocols were performed. No multiplex PCR were executed to allow a better interpretation of the results. Each PCR assay employed a primer pair: *stx1F* (5'-ATAAATCGCCATTCGTTGACTAC-3') and *stx1R* (5'-GAACGCCACTGAGATCATC-3') to amplify a 180 bp fragment of *stx1* gene encoding Shiga toxins 1, *stx2F* (5'-GGCACTGTCTGAAACTGCTCC-3') and *stx2R* (5'-TCGCCAGTTATCTGACATTCTG-3') to amplify a 255 bp fragment of *stx2* gene encoding Shiga toxins 2, *eaeAF* (5'-GACCCGGCACAAGCATAAGC-3') and *eaeAR* (5'-CCACCTGCAGCAACAAGAGG-3') to amplify a 384 bp fragment of *eaeA* gene encoding intimin, and *hlyAF* (5'-GCATCATCAAGCGTACGTTCC-3') and *hlyAR* (5'-AATGAGCCAAGCTGGTTAAGCT-3') to amplify a 534 bp fragment of *hlyA* gene encoding for hemolysin. All primers were previously described by Paton and Paton[20].

Each PCR assay was performed in a volume of 50 µL consisting of 25 µL of EconoTaq PLUS 2x Master Mix (Lucigen Corporation, Middleton, Wisconsin, USA), 0.5 µM of each primer, 3 µL of DNA and distilled water to reach the final volume. The amplifications were carried out in the automated thermal cycler Gene-Amp PCR System 2700 (Perkin Elmer, Norwalk, Connecticut, USA) for 35 cycles. Each cycle consisted of denaturation at 95 °C for 1 min, annealing at 60 °C for 2 min, extension at 72 °C for 1 min; an initial denaturation of 10 min at 95 °C and a final extension of 10 min at 72 °C were done. PCR assays included a reagent blank as negative control. Quantitative Genomic DNA from O157:H7 *E. coli* (ATCC® 43895DQ TM) was included as positive control to assess the protocols, whereas no positive controls were used in the following amplifications to avoid cross-contaminations. PCR products were analyzed by electrophoresis on 1.5% agarose gel at 100 V for 45 min. PCR Sizer 100 bp DNA Ladder (Norgen Biotek, Thorold, Canada) was used as DNA marker.

### 3. Results

The species of 56 hunted and 65 dead birds are shown in Figure 1. All hunted birds were in good nutrition conditions and no lesions were observed during the carcasses' manipulation. The remaining birds were sent to the APS for necropsies, showing fatal traumatic lesions as putative cause of death. No lesions ascribable to infectious and/or parasitic diseases were observed.




Figure 1. Number of examined birds in relation to the avian species.

Table 1. Results of PCR assays for the detection of *E. coli* virulence genes in relation to the animal species.

Species	Genes						Total
	<i>stx1</i>	<i>eaeA</i>	<i>hlyA</i>	<i>eaeA/hlyA</i>	<i>stx1/eaeA</i>	<i>stx1/stx2/eaeA/hlyA</i>	
Seagull ( <i>L. michahellis</i> )	3	5					8
Pigeon ( <i>C. livia</i> )	2	3					5
Mallard duck ( <i>A. platyrhynchos</i> )			1	2			3
Common teal ( <i>A. crecca</i> )	1						1
Grey heron ( <i>A. cinerea</i> )	1						1
Eurasian wigeon ( <i>A. penelope</i> )					1		1
Pintail ( <i>A. acuta</i> )	1						1
Little owl ( <i>A. noctua</i> )						1	1
Total	8	8	1	2	1	1	21

### 4. Discussion

The present study was carried out on several bird species found dead or hunted that lived in different environments. Most of them were animals roaming within an aquatic environment, such as seagulls and waterfowl, whereas others usually live in hinterland such as raptors. Moreover, feral pigeons which reach urban, peri-urban and rural areas were also examined.

The results show that these animals may harbor diarrhea-inducing *E. coli* strains carrying genes for virulence factors able to determine

PCR protocols allowed the amplification and the detection of the four investigated genes fragments when the DNA positive control was tested.

Overall, 21(17.35%) fecal samples resulted positive for at least one of the investigated genes. In detail, 12(9.92%) samples were positive for *eaeA*, 10(8.26%) for *stx1*, 4(3.31%) for *hlyA* and 1(0.83%) for *stx2*.

In particular, 8(6.61%) animals were positive only for *stx1*, 8(6.61%) only for *eaeA*, 1(0.83%) only for *hlyA*, 2(1.65%) for *eaeA* and *hlyA*, 1(0.83%) for *stx1* and *eaeA*, 1(0.83%) for *stx1*, *stx2*, *eaeA* and *hlyA*.

Considering the examined avian species, positive samples were detected in 8 seagulls [*Larus michahellis* (*L. michahellis*)], 5 pigeons [*Columba livia* (*C. livia*)], 3 mallard ducks [*Anas platyrhynchos* (*A. platyrhynchos*)], 1 common teal [*Anas crecca* (*A. crecca*)], 1 grey heron [*Ardea cinerea* (*A. cinerea*)], 1 little owl [*Athene noctua* (*A. noctua*)], 1 Eurasian wigeon [*Anas penelope* (*A. penelope*)] and 1 pintail [*Anas acuta* (*A. acuta*)]. Table 1 reports the distribution of positive animals in relation to the detected genetic profiles. Overall, 17.35% (21/121) birds were positive for the detected genes, of which 4.13% (5/121) were pigeons, 5.79% (7/121) were waterfowl, 6.61% (8/121) were seagulls, and 0.82% (1/121) were raptors.

disease both in animals and humans.

The present investigation found the 8.26% of birds positive for *stx1* gene, alone or in combination with other virulence genes. This result is in contrast with a study recently carried out in Brazil that did not detect avian feces positive for *stx1* gene[6].

Except in the case of one animal, no positive response for *stx2* gene was found. This is a comforting result, considering that *stx-2* toxin is more toxic than *stx-1* and is often associated with HUS in children[21]. Similar results were obtained by Sanches et al.[6], which found 3/401 (0.75%) *stx2*-positive birds.

One bird (*A. platyrhynchos*) was positive only for *hlyA* gene, whereas two others (*A. platyrhynchos*) were positive for both *hlyA* and *eaeA* genes, suggesting they were spreaders of EHEC strains.

The gene *eaeA*, characterizing EPEC strains, was found alone in eight (6.61%) birds, five seagulls and three pigeons, and in combination with *stxI* in one wigeon. This value is in agreement with the study by Sanches *et al.*[6] that found 5.74% of *eaeA*-positive wild birds. Prevalence obtained in other investigations is difficult to compare to our results, since the populations examined are different from our study.

However, other authors mainly found *eaeA*-positive birds among Columbiformes[6,22]. In this study *stx*, *eaeA* and *hlyA* genes were mainly detected in the fecal samples collected from seagulls, waterfowl and pigeons.

Seagulls (*L. michahellis*) are omnivorous aquatic birds, which usually live along the marine coasts. However, they can be found around lakes and in urban/peri-urban areas. They eat mainly fish, rats, animal carcasses, but more often they feed on human food waste found in garbage bins or in landfills[23].

Waterfowl, including ducks and geese, are largely found in wetlands; they indeed can be found in areas with rivers and lakes, but also small ponds, including those presenting in public or private parks. These animals are able to travel large distances per day dispersing pathogens over wide areas. They are largely involved in the epidemiology of some infectious diseases, such as Avian Influenza and Newcastle Disease, because they may excrete/contract the pathogens in the wetland environments shared by numerous birds[24]. At the same way, geese and ducks can contaminate surface water and pasture with other pathogens, including *E. coli*. Moreover, the contaminated surface water, if used for irrigation, represents a further source of infection for people.

Waterfowl are the most common hunted birds. If infected by *E. coli*, they represent a dangerous source of infection for hunters and other people during the carcasses' manipulation, a procedure often carried out in kitchens.

Positive responses detected among feral pigeons (*C. livia*) are relevant for the public health, as well. They are synanthropic birds largely present in urban and peri-urban areas and can contaminate, with their droppings, environments where people are also living[25]. Moreover, pigeons may be source of infection for other animals, such as birds, mammals and reptiles also in rural areas. In particular, pigeons, such as other wild birds, frequently live around farms and are carriers of pathogens for livestock.

During this investigation, a fecal sample collected from an owl (*A. noctua*) scored positive for all the investigated genes. Bacteriological examinations were carried out on this sample and an *E. coli* strain was isolated and typed by cultural and staining characteristics and API 20E System (BioMérieux, Marcy l'Etoile, France). The strain, submitted to molecular analyses, resulted to

have the four investigated virulence genes, suggesting that the animal harbored a STEC strain. On the basis of the available literature, this is the first report of STEC infection in an owl, thus no information about possible pathology due to this type of *E. coli* in owls is known.

Owl is a carnivorous bird of prey that lives mainly on a diet of insects, earthworms, small vertebrates including amphibians, reptiles, birds and small mammals, such as mice, rats, shrews, lagomorphs, voles and moles. This STEC-positive animal could have ingested the pathogen feeding on a prey. This result suggests that STEC strains are circulating, even if not largely, among wildlife, as also demonstrated by other studies that found STEC in Columbiformes[6,22].

In particular, the source of infection for the positive owl could be rodents, which are largely found in rural areas, such as livestock farms. A previous study detected the same STEC strains in feces from rodents and cattle[9].

The obtained results do not clarify whether wild birds can act as spillover hosts or reservoirs for infection. *E. coli* present in the feces of the investigated birds could be transient or resident bacteria in the intestinal tract. Regardless STEC, EPEC and EHEC strains replicate or not in the avian intestine, the presence of these pathogens in the droppings of birds is a severe threat for health of other animals and humans.

## 5. Conclusion

This preliminary investigation, even though carried out on a limited number of samples, shows that wild birds can harbor and excrete in their feces pathogenic *E. coli* strains able to infect and determine disease in other animal species and humans. The highest rates of infection were found among seagulls, waterfowl and feral pigeons, constituting a potential source of *E. coli* for people, companion animals and livestock, as they live in urban, peri-urban and rural environments. Considering that bacterial prophylactic measures on wild birds are not achievable, environmental hygienic actions are basic to reduce the pathogens' spreading. Attention should be paid to the waste management, bins and dumps, which could attract pigeons and gulls near to the human environment. Moreover, surface water should not be used for irrigation and watering of the animals and measures against wild birds are necessary to reduce the environment contaminations with avian droppings.

## Conflict of interest statement

The authors declare no conflict of interest.

## References

- [1] Croxen MA, Finlay BB. Molecular mechanisms of *Escherichia coli* pathogenicity. *Nat Rev Microbiol* 2010; **8**(1): 26-38.
- [2] Kaper JB, Nataro JP, Mobley HL. Pathogenic *Escherichia coli*. *Nat Rev Microbiol* 2004; **2**(2): 123-140.
- [3] Bielaszewska M, Tarr PI, Karch H, Zhang W, Mathys M. Phenotypic and molecular analysis of tellurite resistance among enterohemorrhagic *Escherichia coli* O157:H7 and sorbitol-fermenting O157: NM clinical isolates. *J Clin Microbiol* 2005; **43**(1): 452-4.
- [4] Welinder-Olsson C, Kaijser B. Enterohemorrhagic *Escherichia coli* (EHEC). *Scand J Infect Dis* 2005; **37**(6-7): 405-416.
- [5] Centers for Disease Control and Prevention (CDC). *National Shiga toxin-producing Escherichia coli (STEC) Surveillance Annual Report, 2013*. Atlanta: US Department of Health and Human Services, CDC; 2016.
- [6] Sanches LA, Gomes MDS, Teixeira RHF, Cunha MPV, Oliveira MGX, Vieira MAM, et al. Captive wild birds as reservoirs of enteropathogenic *E. coli* (EPEC) and Shiga-toxin producing *E. coli* (STEC). *Braz J Microbiol* 2017; **48**(4): 760-763.
- [7] Persad AK, LeJeune JT. Animal reservoirs of Shiga toxin-producing *Escherichia coli*. *Microbiol Spectr* 2014; **2**(4): EHEC-0027-2014.
- [8] Wallace JS, Cheasty T, Jones K. Isolation of verocytotoxin producing *Escherichia coli* O157 from wild birds. *J Appl Microbiol* 1997; **82**(3): 399-404.
- [9] Nielsen EM, Skov MN, Madsen JJ, Lodal J, Jespersen JB, Baggesen DL. Verocytotoxin-producing *Escherichia coli* in wild birds and rodents in close proximity to farms. *Appl Environ Microbiol* 2004; **70**(11): 6944-6947.
- [10] Schouten JM, van de Giessen AW, Frankena K, De Jong MCM, Graat EAM. *Escherichia coli* O157 prevalence in Dutch poultry, pig finishing and veal herds and risk factors in Dutch veal herds. *Prev Vet Med* 2005; **70**(1-2): 1-15.
- [11] Wasteson Y, Johannessen GS, Bruheim T, Urdahl AM, O'Sullivan K, Roryik LM. Fluctuations in the occurrence of *Escherichia coli* O157:H7 on a Norwegian farm. *Lett Appl Microbiol* 2005; **40**(5): 373-377.
- [12] Foster G, Evans J, Knight HI, Smith AW, Gunn GJ, Allison LJ, et al. Analysis of feces samples collected from a wild bird garden feeding station in Scotland for the presence of verocytotoxin-producing *Escherichia coli* O157. *Appl Environ Microbiol* 2006; **72**(3): 2265-2267.
- [13] Pedersen K, Clark L. A review of Shiga toxin *Escherichia coli* and *Salmonella enterica* in cattle and free-ranging birds: Potential association and epidemiological links. *Human-Wildlife Conflict* 2007; **1**(1): 68-77.
- [14] Kauffman MD, LeJeune J. European starlings (*Sturnus vulgaris*) challenged with *Escherichia coli* O157 can carry and transmit the human pathogen to cattle. *Lett Appl Microbiol* 2011; **53**(6): 596-601.
- [15] Callaway TR, Edrington TS, Nisbet DJ. Isolation of *Escherichia coli* O157:H7 and *Salmonella* from migratory brown-headed cowbirds (*Molothrus ater*), common Grackles (*Quiscalus quiscula*), and cattle egrets (*Bubulcus ibis*). *Foodborne Pathog Dis* 2014; **11**(10): 791-794.
- [16] Chandran A, Mazumder A. Occurrence of diarrheagenic virulence genes and genetic diversity in *Escherichia coli* isolates from fecal material of various avian hosts in British Columbia, Canada. *Appl Environ Microbiol* 2014; **80**(6): 1933-1940.
- [17] Borges CA, Cardozo MV, Beraldo LG, Oliveira ES, Maluta RP, Barboza KB, et al. Wild birds and urban pigeons as reservoirs for diarrheagenic *Escherichia coli* with zoonotic potential. *J Microbiol* 2017; **55**(5): 344-348.
- [18] Dipineto L, Santaniello A, Fontanella M, Lagos K, Fioretti A, Menna LF. Presence of Shiga toxin-producing *Escherichia coli* in O157:H7 in living layer hens. *Lett Appl Microbiol* 2006; **43**(3): 293-295.
- [19] Santaniello A, Gargiulo A, Borrelli L, Dipineto L, Cuomo A, Sensale M, et al. Survey of Shiga toxin-producing *Escherichia coli* O157:H7 in urban pigeons (*Columba livia*) in the city of Napoli, Italy. *Ital J Anim Sci* 2007; **6**: 313-316.
- [20] Paton AW, Paton JC. Detection and characterization of Shiga toxigenic *Escherichia coli* by using multiplex PCR assays for *stx1*, *stx2*, *eaeA*, enterohemorrhagic *E. coli* hlyA, rfbO111, and rfbO157. *J Clin Microbiol* 1998; **36**(2): 598-602.
- [21] Croxen MA, Law RJ, Scholz R, Keeney KM, Wlodarska M, Finlay BB. Recent advances in understanding enteric pathogenic *Escherichia coli*. *Clin Microbiol Rev* 2013; **26**(4): 822-880.
- [22] Kobayashi H, Kanazaki M, Hata E, Kubo M. Prevalence and characteristics of *eae*- and *stx*-positive strains of *Escherichia coli* from wild birds in the immediate environment of Tokyo Bay. *Appl Environ Microbiol* 2009; **75**: 292-295.
- [23] Maciusik B, Lenda M, Skorka P. Corridors, local food resources, and climatic conditions affect the utilization of the urban environment by the Black-headed Gull *Larus ridibundus* in winter. *Ecol Res* 2010; **25**(2): 263-272.
- [24] Alexander DJ. Orthomyxoviridae: Avian Influenza. In: Pattison M, McMullin PF, Bradbury JM, Alexander D. (eds.) *Poultry diseases*. 6th ed. Edinburgh: Saunders Elsevier; 2008, p. 317-332.
- [25] Haag-Wackernagel D, Moch H. Health hazards posed by feral pigeons. *J Infect* 2004; **48**(4): 307-313.